Growing wildlife populations and trophic rewilding in Sweden plus some insights from African systems

Joris Cromsigt, Megafauna & Sustainability unit

Senior lecturer @ Dept. of Wildlife, Fish & Environmental studies, SLU, Umeå, Sweden

Honorary Professor @ Centre for African Conservation Ecology, Nelson Mandela University, Port Elizabeth, South Africa

Lecturer @ Copernicus Institute of Sustainable Development, Utrecht University, the Netherlands









20 years of experience working in South African systems
Extensive experience in Scandinavian, Dutch and Polish (Białowieża Primeval Forest) systems

The Netherlands: (in)famous for its Rewilding "experiments"









Growing wildlife populations: large changes in Sweden's ungulate community

Relative increase in N of individuals shot (1965-2004)




Expansion of deer species inland and northwards

Bag statistics per Moose Management Area for 2015-2016 hunting year





Beyond Moose program: Consequences of ungulate rewilding













Opportunities for Trophic rewilding in Sweden

Challenge to conserve species that are adapted to open ecosystems















Connected to the loss of extensive farming and land abandonment i.e., biodiversity connected to "anthropogenic habitat"



Review 🙃 Full Access

Farmland abandonment: threat or opportunity for biodiversity conservation? A global review

Cibele Queiroz ⋈, Ruth Beilin, Carl Folke, Regina Lindborg

First published: 01 June 2014 | https://doi.org/10.1890/120348 | Citations: 123

Land abandonment, landscape, and biodiversity: questioning the restorative character of the forest transition in the Mediterranean

Iago Otero 1,2, Joan Marull 3, Enric Tello 4, Giovanna L. Diana 3, Manel Pons 3, Francesc Coll 3 and Martí Boada 2

Ecology and Society 20(2): 7




Authors

Authors and affiliations

Jiří Reif 🔀 , Pavel Marhoul, Oldřich Čížek, Martin Konvička



A African perspective on this






William J. Bond



Challenging traditional view and definition of biomes

Biomes as climate-controlled units (whittaker's diagram)





Climate and soils DO NOT rule the world

Plant consumers (fire, herbivory) control many biomes

Journal of Vegetation Science 16: 261-266, 2005 © IAVS; Opulus Press Uppsala.

261

INVITED PERSPECTIVE

Large parts of the world are brown or black: A different view on the 'Green World' hypothesis


Bond, William J.

Botany Department, University of Cape Town, Private Bag, Rondebosch, 7701, South Africa; Fax +27 216504041; E-mail bond@botzoo.uct.ac.za



William J. Bond







PRINCIPLES OF LIFE, Figure 42.10
© 2012 Sinauer Associates, Inc.



Consumers can strongly control woody biomass





Journal of Vegetation Science 16: 261-266, 2005 © IAVS; Opulus Press Uppsala.

26

INVITED PERSPECTIVE

Large parts of the world are brown or black: A different view on the 'Green World' hypothesis

Bond, William J.

Botany Department, University of Cape Town, Private Bag, Rondebosch, 7701, South Africa; Fax +27 216504041; E-mail bond@botzoo.uct.ac.za





Open Ecosystems: consumer-controlled ecosystems or "ecosystems uncertain"

Open Ecosystems

Fire-driven system

Herbivore-driven system Climate-driven system

Black world




Brown world



Green world







PRINCIPLES OF LIFE, Figure 42.10

© 2012 Sinauer Associates, Inc.



Large extent of the world are 'ecosystems uncertain'!




Figure 5.1 Distribution of "ecosystems uncertain" (dark shading), which are areas categorized as either grasslands or one of the biomes dominated by woody plants in the Whittaker biome diagram (reprinted with permission from Bond, 2005).



What does this mean in terms of rewilding?

- Well ... calls for a paradigm shift
- Instead of restoring European open habitat specialists through restoring "anthropogenic habitats and drivers"
- We need to refocus on restoring Europe's Open Ecosystems through restoring its drivers -> fire and herbivory
 - i.e., Trophic Rewilding




Restore open ecosystems by restoring the role of consumers







Restoring functionality in Europe's large herbivore community (large grazers)






Rewilding with free-ranging cattle and horses

As substitutes of extinct Aurochs and Tarpan

Aurochs went extinct in 1627



Tarpan went extinct in 1879




Restoring Open Ecosystems through Trophic Rewilding = restoring herbivory AND fire





We need out-of-the-box experiments to start testing these ideas





Fire and grazing experiment, Nordens Ark, Sweden

- Karin Amsten¹ (PhD student)
- Mats Niklasson¹ (main supervisor)
- Jenny Loberg¹
- Joris Cromsigt
- Marcin Churski²
- Dries Kuijper²



¹Nordens Ark Zoo & SLU

²Mammal Research Institute, Bialowieza, Poland








Experiment replicated in 24 sites across Nordens Ark's Ekopark



Survival after three years (2 spring burning events)



Amsten et al. Submitted to New Phytologist



Interaction between grazing and fire



SLL


Grazing treatment hardly burnt





Lots to learn still...

- Particularly from science coming out of southern hemisphere savannas and grasslands
- This can guide efforts to restore European Open Ecosystems through trophic rewilding





"Rewilding our minds": rethink some of our terminology?


13th Eurasian Grassland Conference

Management and Conservation of Semi-natural grasslands. from theory to practice
Sighişoara, Romania
20-24 September 2016



Citation from Otero et al. 2015. *Ecology and Society* 20:7. "We (1) question the alleged relationship between land abandonment and ecosystem recovery; (2) show that the assumed restorative character of the forest transition is based on the underestimation of the ecological importance of nonforest habitats"





PRINCIPLES OF LIFE, Figure 42.10

© 2012 Sinauer Associates, Inc.

